

The Risk Quotient Method is the means used by EFED to integrate the results of exposure and ecotoxicity data. For this method, risk quotients (RQs) are calculated by dividing exposure estimates by ecotoxicity values (*i.e.*, $RQ = EXPOSURE/TOXICITY$), for both acute and chronic endpoints. Acute RQs use the LD₅₀ or LC₅₀ as the toxicity value, and chronic RQs use the NOAEC or NOAEL as the toxicity value. The RQs are then compared to OPP's levels of concern (LOCs). These LOCs are criteria used by OPP to indicate potential risk to non-target organisms and the need to consider regulatory action. EFED has defined LOCs for acute risk, potential restricted use classification, and for endangered species.

An exceedence of the criteria indicate that a pesticide used as directed has the potential to cause adverse effects on non-target organisms. LOCs currently address the following risk presumption categories:

- (1) acute - there is a potential for acute risk; regulatory action may be warranted in addition to restricted use classification;
- (2) acute restricted use - the potential for acute risk is high, but this may be mitigated through restricted use classification
- (3) acute endangered species - the potential for acute risk to endangered species is high, regulatory action may be warranted, and
- (4) chronic risk - the potential for chronic risk is high, regulatory action may be warranted.

Currently, EFED does not perform assessments for chronic risk to plants, acute or chronic risks to non-target insects, or chronic risk from granular/bait formulations to mammalian or avian species.

The ecotoxicity test values (*i.e.*, measurement endpoints) used in the acute and chronic risk quotients are derived from required studies submitted by the registrant. Test values may also be derived from open literature data included in the ECOTOX database.

Examples of ecotoxicity values derived from short-term laboratory studies that assess acute effects are: (1) LC₅₀ (fish and birds), (2) LD₅₀ (birds and mammals), (3) EC₅₀ (aquatic plants and aquatic invertebrates), and (4) EC₂₅ (terrestrial plants). Examples of toxicity test effect levels derived from the results of long-term laboratory studies that assess chronic effects are: (1) LOAEL (birds, fish, and aquatic invertebrates), and (2) NOAEL (birds, fish and aquatic invertebrates). The NOAEL is generally used as the ecotoxicity test value in assessing chronic effects.

Risk presumptions, along with the corresponding RQs and LOCs are summarized in Table G-1.

Table G-1: Risk Presumptions and LOCs

Risk Presumption	RQ	LOC
Birds¹		
Acute Risk	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day	0.5
Acute Restricted Use	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day (or LD ₅₀ < 50 mg/kg)	0.2
Acute Endangered Species	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day	0.1
Chronic Risk	EEC/NOAEC	1
Wild Mammals¹		
Acute Risk	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day	0.5
Acute Restricted Use	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day (or LD ₅₀ < 50 mg/kg)	0.2
Acute Endangered Species	EEC/LC ₅₀ or LD ₅₀ /sqft or LD ₅₀ /day	0.1
Chronic Risk	EEC/NOAEC	1
Aquatic Animals²		
Acute Risk	EEC/LC ₅₀ or EC ₅₀	0.5
Acute Restricted Use	EEC/LC ₅₀ or EC ₅₀	0.1
Acute Endangered Species	EEC/LC ₅₀ or EC ₅₀	0.05
Chronic Risk	EEC/NOAEC	1
Terrestrial and Semi-Aquatic Plants		
Acute Risk	EEC/EC ₂₅	1
Acute Endangered Species	EEC/EC ₀₅ or NOAEC	1
Aquatic Plants²		
Acute Risk	EEC/EC ₅₀	1
Acute Endangered Species	EEC/EC ₀₅ or NOAEC	1

¹ LD₅₀/sqft = (mg/sqft) / (LD₅₀ * wt. of animal)

LD₅₀/day = (mg of toxicant consumed/day) / (LD₅₀ * wt. of animal)

² EEC = (ppm or ppb) in water