


Command® 3ME

Herbicide

For Agricultural or Commercial Use Only

EPA REG. No. 279-3158



EMERGENCY CALLS: 800-331-3148

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

CROP	WEEDS																
Peppers	<p>The following annual Broadleaf weeds and grasses will be controlled or suppressed for 3-5 weeks:</p> <table border="0"> <tr> <td>Barnyardgrass</td> <td>Velvetleaf</td> </tr> <tr> <td>Broadleaf Signalgrass</td> <td>Common Ragweed</td> </tr> <tr> <td>Crabgrass</td> <td>Jimsonweed</td> </tr> <tr> <td>(Large, Smooth)</td> <td>Lambsquarters</td> </tr> <tr> <td>Field Sandbur</td> <td>Prickly Sida</td> </tr> <tr> <td>Foxtail</td> <td>Pennsylvania Smartweed</td> </tr> <tr> <td>(Giant, Green, Robust)</td> <td>Purslane</td> </tr> <tr> <td>Fall Panicum</td> <td>Venice Mallow</td> </tr> </table>	Barnyardgrass	Velvetleaf	Broadleaf Signalgrass	Common Ragweed	Crabgrass	Jimsonweed	(Large, Smooth)	Lambsquarters	Field Sandbur	Prickly Sida	Foxtail	Pennsylvania Smartweed	(Giant, Green, Robust)	Purslane	Fall Panicum	Venice Mallow
Barnyardgrass	Velvetleaf																
Broadleaf Signalgrass	Common Ragweed																
Crabgrass	Jimsonweed																
(Large, Smooth)	Lambsquarters																
Field Sandbur	Prickly Sida																
Foxtail	Pennsylvania Smartweed																
(Giant, Green, Robust)	Purslane																
Fall Panicum	Venice Mallow																

DIRECTIONS FOR USE:

For the suppression of annual grasses and broadleaf weeds in peppers apply broadcast preemergence to weeds, and prior to transplant at 0.67 to 2.67 pints per acre. Use the lower recommended rate on coarse soils and the higher recommended rate on fine soils.

Do not use on banana peppers.

NOTE: Prior to use, refer to the Precautions and Restrictions sections of this document.

SPECIAL PRECAUTIONS:

Off-site movement spray drift or vapors of Command herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and applications on this and the Federal label.

Recommendations for use of this product are based on tests believed to be reliable. However, not all cultivars of all commodities have been tested. Differences in sensitivity have been observed between cultivars. Therefore, temporary whitening and/or yellowing of the treated crop may occur. The crop should grow through this with no adverse impacts.

(Continued next page)

RESTRICTIONS:

Do not make more than one application.

Refer to the SOYBEANS section of the Command 3ME label for replanting instructions. Do not replant treated fields with any crop inconsistent with the Command 3ME label.

FMC Corporation intends that this section 24(c) label be distributed only to end users and/or growers (and/or applicators acting on behalf of growers) who agree to the terms and conditions below, and agree in writing to a waiver and release from all liability by the user and/or grower of FMC for failure to perform and crop damage from use of Command 3ME Herbicide in single preplant incorporated or preemergence ground treatments on peppers. If such terms and conditions are unacceptable, FMC requests the return of Command 3ME Herbicide at once unopened. FMC considers the user to have accepted such terms and conditions upon use of this product.

This product when used single preplant or preemergence ground treatments on peppers may lead to crop injury, loss, or damage. FMC recommends that the user and/or grower test this product in order to determine its suitability for such intended use. FMC makes this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use this product must be made by each individual user and/or grower on the basis of possible crop injury from Command 3ME Herbicide, the severity of weed infestations, the cost of alternative weed controls, and other factors. Because of the risk of crop damage all such use is at the user's and/or grower's risk.

APPLICATION PRECAUTIONS:

Do not apply Command 3ME herbicide within 1,200 feet of the areas listed below:

Towns and Housing Developments

Commercial Vegetable Production (except peppers, pumpkins, succulent peas, sweet corn, sweet potato and winter squash)

Commercial Fruit / Nut Production

Commercial Nurseries

Commercial Greenhouses

Caution must be taken to minimize spray drift as off-site movement of Command can cause foliar whitening or yellowing of some plants. Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Leave an adequate buffer zone between the area to be treated and desirable plants.

Follow the rotational crop restrictions for soybeans (Northern Area) on the federal label.

Command may be tank mixed with labeled amounts of other registered herbicides.

Consult individual labels for use, directions and precautions.

Do not use this product through any type of irrigation system.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED. PERSONAL PROTECTIVE EQUIPMENT AND REENTRY INTERVALS ON THIS 24C OR ON THE BASIC LABEL MUST BE FOLLOWED. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

24(c) Registrant:

FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia, PA 19103

OH000003

The logo for FMC Corporation, consisting of the letters 'FMC' in a bold, sans-serif font. The 'F' and 'M' are connected, and the 'C' is separate. A registered trademark symbol (®) is located to the right of the 'C'.

Command and **FMC** - FMC Trademarks
©2000 FMC Corporation

Code 618 5/2000